

PAPER • OPEN ACCESS

Inhibition of *Listeria monocytogenes* by natural antimicrobial

To cite this article: T Budiati *et al* 2020 *IOP Conf. Ser.: Earth Environ. Sci.* **411** 012042

View the [article online](#) for updates and enhancements.

You may also like

- [Antimicrobial effects and mechanism of plasma activated fine droplets produced from arc discharge plasma on planktonic *Listeria monocytogenes* and *Escherichia coli* O157:H7](#)
K H Baek, H I Yong, J H Yoo *et al.*
- [Immunomagnetic Bead Separation Coupled with a Dithiobis-Succinimidy Propionate \(DSP\)-Modified Immunosensor to Detect *Listeria monocytogenes* in Chicken Skin](#)
Mi-Kyung Park and Jun-Hyun Oh
- [Detection of *Listeria monocytogenes* from clinical specimens](#)
Maitham Ghaly Yousif and Ataa Khalil AL-Shamari



244th Electrochemical Society Meeting

October 8 – 12, 2023 • Gothenburg, Sweden

50 symposia in electrochemistry & solid state science

Abstract submission deadline:
April 7, 2023

Read the call for papers &

submit your abstract!

Inhibition of *Listeria monocytogenes* by natural antimicrobial

T Budiati*¹, Y Wibisono¹, R A Pambayun², M F Fahrezy², R Ariyani², E Kurniawati¹, W Suryaningsih¹, S O N Yudiastuti¹ and A Bakri²

1 Food Engineering Department, Politeknik Negeri Jember, Mastrip PO Box 164 Jember 68101 East Java, Indonesia

2 Food Industrial Technology Department, Politeknik Negeri Jember, Mastrip PO Box 164 Jember 68101 East Java, Indonesia

Email : titik_budiati@polije.ac.id

Abstract. The aim of the study was to evaluate antimicrobial activity of essential oil from plants as natural antimicrobial to inhibit *Listeria monocytogenes*. A total of 6 essential oil extracted from galangal (*Alpinia galanga*), ginger (*Zingiber officinale*), lemongrass (*Cymbopogon citratus*), orange skin (*Citrus sinensis*), rosemary (*Rosmarinus officinalis*) and turmeric (*Curcuma longa*). By using Minimum Inhibitory Concentration (MIC), the highest and lowest antimicrobial activity of essential oil to inhibit the growth of *L. monocytogenes* was found on lemongrass oil ($0.32 \% \pm 0.12 \%$) and turmeric ($7.46 \% \pm 2.79\%$), respectively. By using disc diffusion assay, the highest antimicrobial activity to against *L. monocytogenes* was shown by lemongrass oil (7.46 ± 2.79 mm). There is no antimicrobial activity observed in orange peel oil. Out of the essential oil tested, lemongrass oil showed the most promising natural antimicrobial to inhibit *L. monocytogenes*.

1. Introduction

Essential oils, the natural antimicrobial, have been known for decades and, with increasing demand from changes in food consumer trends and isolation of antibiotic resistant microorganisms, being other to chemical based antimicrobial which introduce the accumulative residue and give an adverse human health. Essential oil are composed by bioactive compounds which known as generally recognized as safe (GRAS) to food and might be required by food industries for ensuring food safety. Xie *et al.* [1] revealed that the biological agents of essential oils including phenolics and polyphenols have to be high activity against pathogenic bacteria [1,2].

L. monocytogenes is a Gram-positive bacteria and become the main causal agent of listeriosis [3,4], the rare but lethal food-borne disease, especially for susceptible group [5]. This group was categorized as young, old, pregnant, and immunocompromised (YOPI) [6]. The contamination of *L. monocytogenes* may occur during food chain and could be spread to food processing equipments and may contaminate food products [7,8].

The study of essential oils, natural antimicrobials, extracted from Indonesia plants has to be investigated further. To do so, this study was aiming to evaluate antimicrobial activity of essential oil from plants as natural antimicrobial to inhibit *Listeria monocytogenes*. Nevertheless, it becomes a need to get the effective antimicrobial and safe to human health.



2. Material and methods

2.1. Essential oil

Six essential oils namely lemongrass (*Cymbopogon citratus*), ginger (*Zingiber officinale*), galangal (*Alpinia galanga*), orange peel (*Citrus sinensis*), rosemary (*Rosmarinus officinalis*) and turmeric (*Curcuma longa*). The pure essential oils were obtained from Happy Green (Indonesia).

2.2. The strains of microorganism

Pure culture of *L.monocytogenes* (Nusaroma, Gajah Mada University, Yogyakarta) was cultured on Agar Listeria Ottavani and Agosti (ALOA) (Himedia, India) with supplement of ALOA (Himedia, India) for 24 until 48 hours at 37°C. The colonies of *L. monocytogenes* were appeared as blue colonies with halo surround the colonies. By using streaking method, a single colony was purified onto culture media namely tryptic soya agar (TSA) (Himedia, India).

2.3 Minimum Inhibitory Concentration (MIC)

MIC was measured as done by Budiati *et al.* [2]. Cultures were grown at 37 °C for 16 hours in Luria Broth (LB, Himedia, India) in shaking incubator. The cultures were diluted and compared the turbidity. This has to be set at 0.5 of McFarland scale. A serial concentration was prepared on Mueller Hinton Broth (Oxoid, United Kingdom). A total of 20 µL of the aliquot was artificially contaminated to the serial dilution. These were incubated at a temperature of 37 °C for 24 hours. MIC value (%) describes the lowest concentration of the bacteria might grow. These values are calculated as mean values of three measurements. The experiment was done in triplicate. A 15 µl aliquot of 10% DMSO was dropped onto a serial dilution as negative control. Penicillin G (10 unit/ml) was used as positive control.

2.4 Disc diffusion assay

By using disc diffusion assay, the antimicrobial activity was measured as the method of Bauer [9]. Cultures were grown at 37 °C for 16 hours in Luria Broth (LB, Himedia, India) in shaking incubator. The cultures were diluted and the turbidity was set at 0.5 of McFarland scale. By using a sterilized cotton swab, the cultures were swabbed onto Mueller Hinton Agar (Oxoid, United Kingdom) and were let about 30 min. Twenty microliter essential oil was dropped on sterilized disc. This disc was then placed onto the agar. This was incubated for 18–24 h at 37 °C. The inhibition zones was measured in four different directions after the incubation period. This was recorded as milimeter. The experiment was done in triplicate. Fifteen microliter aliquot of 10% DMSO was dropped onto a sterile paper disc as negative control. Penicillin G (10 unit/ml) was used as positive control.

2.5 Statistical Analysis

By using one-way ANOVA (SPSS version 13.0), the differences of antimicrobial activity among essential oils to inhibit the growth of *L. monocytogenes* were determined at a significance level of $P < 0.05$.

3. Results and Discussion

Wide-spectrum of antimicrobial extracted from plant namely essential oils to against Gram-positive bacteria have been well documented [10,11]. *L. monocytogenes* is a Gram-positive bacteria that can be inhibited by essential oil [12,13,14]. By using disc diffusion assay (inhibition zone) and MIC, the antimicrobial activity of six different essential oils extracted from plant in Indonesia to against *L. monocytogenes* was studied. This present study found that the highest antimicrobial activity to inhibit *L. monocytogenes* was observed on lemongrass as shown in table 1. This is similar to other study that reported that lemongrass was effective to inhibit Gram-positive bacteria such as *L. monocytogenes* [15].

Table 1. Inhibition zone and Minimum Inhibitory Concentration (MIC) of food-borne pathogenic bacteria *L.monocytogenes* on six different essential oils.

| Essential oil | Inhibition zone (mm) | MIC (%) |
|---------------|--------------------------|--------------------------|
| Lemongrass | 7,46 ± 2,79 ^b | 0,32 ± 0,12 ^a |
| Ginger | 1,11 ± 0,14 ^a | 1,56 ± 0 ^c |
| Galangal | 0,67 ± 0,60 ^a | 2,60 ± 0,01 ^b |
| Orange peel | nd | 3.13 ± 0 ^d |
| Rosemary | 1,29 ± 0,58 ^a | 3.13 ± 0 ^d |
| Turmeric | 0,58 ± 0,52 ^a | 12,5 ± 0 ^e |

Note : ^{a,b} = different alphabet means significant different at P<0.05 in the same column
nd = not detected

L. monocytogenes was composed by peptidoglycan which was disrupted due to the exposure of citrate and other terpene in lemongrass. The devastating effect in the microbial cell by disrupting the cell membrane integrity by this compound may inhibit the growth of *L. monocytogenes* [16, 17, 18]. This may disrupt to the fatty acid biosynthesis and peptidoglycan biosynthesis [19]. Hadjilouka *et al.* [19] also reported that the gene expression in *L. monocytogenes* was observed due to the exposure of lemongrass essential oil. The genes were *accP*, *accA* and *fapR* that involved in fatty acid biosynthesis. Moreover the expression genes of *murR* and *pbpB* was also reported to be downregulated [19] to cause the disruption of peptidoglycan biosynthesis.

The composition of lemongrass was majority composed by geraniol and neral [19]. The other compounds were also reported by Hadjiloka [19]. These were camphene, limonene, γ -cadinene, geraniol, 4-nonanone, isogeraniol, α -pinene, citronellal, eucalyptol, sabinene, isoneral, caryophyllene oxide, geranyl acetate, isoeugenol, 6-methyl-5-hepten-2-one, β -Pinene, myrcene, linalool, (ω)- β -ocimene, chrysanthemal, (ϵ)-caryophyllene, α -Pinene epoxide, δ -Cadinene, (ϵ)- β -Ocimene, isborneol, α -Terpineol, decanal, geranyl formate, geranyl butyrate, rose furan oxide, trans-piperitol, (ϵ)- γ -Bisabolene, tricyclene, epichroital, β -Elemene. The compounds may introduce the downregulation of *hly* and *inlJ* [19]. Those genes played an important role for the interaction between pathogenic bacteria with the host during infectious process [20].

This present study observed that the MIC value of ginger oil, galangal oil, orange peel oil, rosemary oil and turmeric oil were significant different for each other (table 1). However, inhibition zone of ginger oil, galangal oil, orange peel oil, rosemary oil and turmeric oil were not significant different. It indicated that all those essential oil were not effective to inhibit *L. monocytogenes* compared to lemongrass oil. The lowest antimicrobial activity was observed on turmeric oil by using MIC. The majority compound of turmeric was curcumin and other compounds such as camphor, terpenes, lactones, alkaloids, and phenols [21]. The presence of those compounds indicated to be not effective to against *L. monocytogenes*. This is similar to other studies. Thongson *et al.* [22] reported that turmeric had no antimicrobial effect against *L. monocytogenes*. There is no antimicrobial activity was observed on orange peel oil. The compound of orange peel oil was dominated by limonene [23]. In contrary, the study of Fisher and Philips [23] reported that orange peel had antimicrobial effect to *L. monocytogenes*.

4. Conclusion

Lemongrass oil showed the highest antimicrobial effect to against the growth of *L. monocytogenes*. These essential oil seem to be a potential natural antimicrobial for inhibiting *L. monocytogenes* which was safe to food product. This was also a potential natural antimicrobial that could be used as disinfectant to clean food processing equipments which was vulnerable for the growth of *L. monocytogenes* as biofilms.

Acknowledgment

The financial support was provided by PNPB research scheme - Politeknik Negeri Jember is gratefully acknowledged.

References

- [1] Xie Y, Yang W, Tang F, Chen X and Ren L 2015 Antibacterial activities of flavonoids: structure-activity relationship and mechanism *Current Medicinal Chemistry* **22** 132-149
- [2] Budiati T, Suryaningsih W, Umaroh S, Poerwanto B, Bakri A and Kurniawati E 2018 Antimicrobial activity of essential oil from Indonesian medicinal plants against food-borne pathogens *IOP Conference Series: Earth and Environmental Science* **207** 012036
- [3] Buchanan R L, Gorris L G, Hayman M M, Jackson T C and Whiting R C 2017 A review of *Listeria monocytogenes*: an update on outbreaks, virulence, dose-response, ecology, and risk assessments *Food control* **75** 1-13
- [4] Budiati T, Rusul G, Wan-Abdullah W N, Arip Y M and Ahmad R 2013 Comparison of sample preparation for the isolation of *Listeria* species in naturally contaminated catfish and tilapia samples *Emirates Journal of Food and Agriculture* **25** 613-617
- [5] Madjunkov M, Chaudhry S and Ito S 2017 Listeriosis during pregnancy *Archives of gynecology and obstetrics* **296** 143-152
- [6] Cossu F, Spanu C, Deidda S, Mura E, Casti D, Pala C, Lamon S, Spanu V, Ibba M, Marrocu E and Scarano C 2016 *Listeria* spp. and *Listeria monocytogenes* contamination in ready-to-eat sandwiches collected from vending machines *Italian journal of food safety* **5**
- [7] Camargo A C, Woodward J J, Call D R and Nero L A 2017 *Listeria monocytogenes* in food-processing facilities, food contamination, and human listeriosis: the Brazilian scenario *Foodborne Pathogens and Disease* **14** 623-636
- [8] Titik B., Naiyana C, 2010 Exposure Assessment for *Listeria monocytogenes* in cooked ham *International Food Research Journal* **17** 267-280
- [9] Bauer A, Kirby W, Sherris J and Turck M 1966 Antibiotic susceptibility testing by a standardized single disc method *Am. J. Clin. Pathol.* **45** 493
- [10] de Oliveira M M M, Brugnera D F, Cardoso M, Alves G E and Piccoli R H 2010 Disinfectant action of *Cymbopogon* sp. essential oils in different phases of biofilm formation by *Listeria monocytogenes* on stainless steel surface *Food Control* **21** 549-553
- [11] Van Vuuren S F 2008 Antimicrobial activity of South African medicinal plants. *Journal of ethnopharmacology* **119** 462-472
- [12] Pesavento G, Calonico C, Bilia A R, Barnabei M, Calesini F, Addona R, Mencarelli L, Carmagnini L, Di Martino M C and Nostro AL 2015 Antibacterial activity of Oregano, Rosmarinus and Thymus essential oils against *Staphylococcus aureus* and *Listeria monocytogenes* in beef meatballs *Food Control* **54** 188-199
- [13] de Medeiros B, da Costa M I, de Oliveira J A, Gomes-Neto K Á R, Tavares N J, Magnani J F M and de Souza E L 2016. Efficacy of the combined application of oregano and rosemary essential oils for the control of *Escherichia coli*, *Listeria monocytogenes* and *Salmonella* Enteritidis in leafy vegetables *Food control* **59** 468-477
- [14] Lee G, Kim Y, Kim H, Beuchat L R and Ryu J H 2018. Antimicrobial activities of gaseous essential oils against *Listeria monocytogenes* on a laboratory medium and radish sprouts. *International journal of food microbiology* **265** 49-54
- [15] Barbosa L N, Rall V L M, Fernandes A A H, Ushimaru P I, da Silva P I and Fernandes Jr A 2009 Essential oils against foodborne pathogens and spoilage bacteria in minced meat. *Foodborne pathogens and disease*, **6** 725-728
- [16] Somolinos M, García D, Pagán R and Mackey B M 2008 Relationship between 606 sublethal injury and microbial inactivation by the combination of high hydrostatic pressure and citral or tert-butyl hydroquinone. *Applied and Environmental Microbiology* **74** 7570-7577

- [17] Somolinos M, Garcia D, Condon S, Mackey B and Pagan R 2009 Inactivation of *Escherichia coli* by citral *Journal of Applied Microbiology* **108** 1928–1939
- [18] Park M J, Gwak K S, Yang I, Kim K W, Jeung, E B, Chang J W, Choi I G 2009 Effect of citral, eugenol, nerolidol, and α -terpineol on the ultrastructural changes of Trichophyton mentagrophytes. *Fitoterapia* **80** 290–296
- [19] Hadjilouka A, Mavrogiannis G, Mallouchos A, Paramithiotis S, Mataragas M and Drosinos E H 2017 Effect of lemongrass essential oil on *Listeria monocytogenes* gene expression. *LWT* **77** 510-516
- [20] Bierne H, Sabet C, Personnic N and Cossart P 2007 Internalins: a complex family of leucine-rich repeat-containing proteins in *Listeria monocytogenes* *Microbes and Infection* **9** 1156-1166
- [21] Mukunthan K S, Balaji B and Patel T N 2018 Black turmeric database: A database of natural compounds from curcuma caesia roxb *Asian Journal of Pharmaceutical and Clinical Research* **11** 406-408
- [22] Thongson C, Davidson P M, Mahakarnchanakul W and Vibulsresth P 2005 Antimicrobial effect of Thai spices against *Listeria monocytogenes* and *Salmonella* Typhimurium DT104 *Journal of Food Protection* **68** 2054-2058
- [23] Fisher K and Phillips C A 2006. The effect of lemon, orange and bergamot essential oils and their components on the survival of *Campylobacter jejuni*, *Escherichia coli* O157, *Listeria monocytogenes*, *Bacillus cereus* and *Staphylococcus aureus* in vitro and in food systems *Journal of Applied Microbiology* **101** 1232-1240