

# Gayuh Syaikhullah

## Exploring the Relationship Between Growth Hormone Secretagogue Receptor (GHSR) Gene and Body Proportions in ...

 Quick Submit

 Quick Submit

 Politeknik Negeri Jember

---

### Document Details

Submission ID

trn:oid::1:2980502861

Submission Date

Aug 9, 2024, 1:22 PM GMT+7

Download Date

Aug 9, 2024, 1:31 PM GMT+7

File Name

8\_Syaikhullah.pdf

File Size

536.3 KB

8 Pages

3,943 Words

20,420 Characters

# 17% Overall Similarity

The combined total of all matches, including overlapping sources, for each database.





## Filtered from the Report

- ▶ Bibliography
- ▶ Quoted Text
- ▶ Cited Text
- ▶ Small Matches (less than 8 words)




## Exclusions

- ▶ 7 Excluded Matches

## Match Groups

-  **42 Not Cited or Quoted 17%**  
Matches with neither in-text citation nor quotation marks
-  **0 Missing Quotations 0%**  
Matches that are still very similar to source material
-  **0 Missing Citation 0%**  
Matches that have quotation marks, but no in-text citation
-  **0 Cited and Quoted 0%**  
Matches with in-text citation present, but no quotation marks

## Top Sources

- 15%  Internet sources
- 9%  Publications
- 2%  Submitted works (Student Papers)

## Integrity Flags

### 0 Integrity Flags for Review

No suspicious text manipulations found.

Our system's algorithms look deeply at a document for any inconsistencies that would set it apart from a normal submission. If we notice something strange, we flag it for you to review.

A Flag is not necessarily an indicator of a problem. However, we'd recommend you focus your attention there for further review.

### Match Groups

- 42 Not Cited or Quoted 17%**  
Matches with neither in-text citation nor quotation marks
- 0 Missing Quotations 0%**  
Matches that are still very similar to source material
- 0 Missing Citation 0%**  
Matches that have quotation marks, but no in-text citation
- 0 Cited and Quoted 0%**  
Matches with in-text citation present, but no quotation marks

### Top Sources

- 15% Internet sources
- 9% Publications
- 2% Submitted works (Student Papers)

### Top Sources

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

<b>1</b>	Internet	discovery.researcher.life	4%
<b>2</b>	Internet	core.ac.uk	2%
<b>3</b>	Student papers	UC, San Diego	2%
<b>4</b>	Internet	media.neliti.com	1%
<b>5</b>	Internet	ijsrset.com	1%
<b>6</b>	Internet	ojs.uho.ac.id	1%
<b>7</b>	Internet	academic.oup.com	0%
<b>8</b>	Internet	www.e3s-conferences.org	0%
<b>9</b>	Publication	W L N Aliyya, R R Noor, Jakaria. "Exploring SNPs (Single Nucleotide Polymorphis..."	0%
<b>10</b>	Internet	www.veterinaryworld.org	0%

11	Publication	Jing Ni, Feng You, Jianhe Xu, Dongdong Xu, Aiyun Wen, Zhihao Wu, Yongli Xu, Peij...	0%
12	Internet	link.springer.com	0%
13	Publication	Kholifah Kholifah, Puji Hartati, Dewi Pranatasari. " The Effect of Vitamin E and Sel...	0%
14	Publication	Wenya Li, Danli Liu, Shuqi Tang, Donghua Li et al. "A multiallelic indel in the prom...	0%
15	Internet	icgtd.ucr.edu	0%
16	Internet	jurnal.ipb.ac.id	0%
17	Publication	M M D Utami, D Pantaya, A Agus. "Addition of Garlic Extract in Ration to Reduce C...	0%
18	Publication	Toivo Jurimae, Neil Armstrong, Jaak Jurimae. "Children and Exercise XXIV - The Pr...	0%
19	Internet	www.tandfonline.com	0%
20	Publication	Aan Awaludin, Raden Wisnu Nurcahyo, Dwi Priyowidodo, Yudhi Ratna Nugraheni,...	0%
21	Publication	D Pantaya, S Wulandari, A P Yulinarsari, H Poernomo. "Evaluation of rubber seed ...	0%
22	Publication	G. C. Everitt. "Effects of sex and gonadectomy on the growth and development of ...	0%
23	Internet	businessdocbox.com	0%
24	Internet	docslib.org	0%

25	Internet	
ejrr.gau.ac.ir		0%
<hr/>		
26	Internet	
eprints.whiterose.ac.uk		0%
<hr/>		
27	Internet	
seminar.fpp.undip.ac.id		0%

# Exploring the Relationship Between Growth Hormone Secretagogue Receptor (GHSR) Gene and Body Proportions in Bangkok Chickens: Insights from DNA Sequencing and Hin6I Enzyme-Restricted PCR-RFLP Analysis

Gayuh Syaikhullah<sup>1\*</sup>, Rosa Tri Hertamawati<sup>1</sup>, Muhammad Adhyatma<sup>2</sup>

<sup>1</sup>Poultry Business Management Study Program, Department of Animal Husbandry, Politeknik Negeri Jember, Jember, Indonesia, <sup>2</sup>Animal Production Study Program, Department of Animal Husbandry, Politeknik Negeri Jember, Jember, Indonesia.

\*Corresponding author: [gayuh\\_syaikhullah@polije.ac.id](mailto:gayuh_syaikhullah@polije.ac.id)

## Abstract

The purpose of this study was to determine the diversity of the growth hormone secretagogue receptor (GHSR) gene and also to analyze the association of the GHSR gene with the body proportions of Bangkok chickens. A total of 125 Bangkok chickens were reared, and blood samples were taken. Bangkok chickens were kept in colony cages with ad libitum feeding and drinking. The GHSR gene polymorphism was determined using the PCR-RFLP method. The restriction enzyme used in this study was Hin6I. The data analyzed were indicators of body proportions, which included data on body weight, carcass weight, and commercial cut weight. Single nucleotide polymorphism (SNP) identification using the Molecular Evolutionary Genetics Analysis 7 program with reference number AB095994.1. Allele frequency values, genotype frequencies, and Hardy-Weinberg balance values were also analyzed. The association between the GHSR gene and the traits observed in Bangkok chickens was analyzed using the T-test. The results of the study showed that the GHSR gene in Bangkok chickens had two genotypes, namely TT and CT. The values for  $H_o$  and  $H_e$  were 0.224 and 0.198, respectively. The genotype frequencies of TT and CT were 0.776 and 0.224, respectively. The two genotypes were associated with body weight, carcass weight, and commercial weight ( $p < 0.05$ ). For all measured criteria, the TT genotype showed a higher weight value than the CT genotype. The GHSR gene has the potential to be used as a genetic marker for the selection process on body weight, carcass weight, and commercial weight traits.

Keywords: GHSR gene, Bangkok chicken, PCR, RFLP

Received: 30 August 2023

Revised: 15 January 2024

Accepted: 16 February 2024

## INTRODUCTION

Indonesia has a large number of local chicken livestock with significant development potential. At least 34 local chicken species with various morphological traits can be found in Indonesia (Ningsih *et al.*, 2022). Indonesia boasts a diverse array of local chicken breeds, encompassing both original varieties and those modified over decades or even centuries. Among them is the Bangkok Chicken, a descendant of Thailand's regional chicken, historically known as the ancestor of the King's chicken. Having been cultivated for an extensive period and widely adopted by Indonesians since 1960, the Bangkok Chicken is officially recognized as a local Indonesian breed. Its popularity stems from

numerous advantages, such as a robust body structure and exceptional endurance, making it a preferred choice for various applications. (Badaruddin *et al.*, 2017). In addition to that, Bangkok chickens possess high economic value. It is not surprising that breeders maintain them for business purposes, rather than merely as a hobby and source of pride. Another notable attribute of Bangkok chickens is the exceptionally good quality of their sperm (Hijriyanto, 2017). This quality promote Bangkok chickens suitable as male candidates in the process of establishing new superior local chicken strains.

Despite these advantages, Bangkok chickens also have disadvantages, such as a relatively slow growth rate. Almost all types of local Indonesian chickens tend to share this drawback, exhibiting

low productivity that falls below their genetic potential (Nataamijaya, 2010). Muryanto *et al.* (2002) stated that the development of local chickens has constraints on the slow rate of growth. Therefore, it is necessary to improve the genetic quality of Bangkok chickens to increase the efficiency of this type of chicken cultivation. One method that can be used is selection based on specific genes that affect growth traits.

The growth hormone secretagogue receptor (GHSR), or growth hormone secretagogue receptor, is involved in many physiological functions, including pituitary growth hormone secretion, food intake, and energy expenditure (Fang *et al.*, 2010). The GHSR gene was identified as being associated with growth performance in kampung chickens (Syaikhullah *et al.*, 2017). The GHSR gene is a gene that plays an important role as the main receptor in the process of GH secretion, or what is often called growth hormone (Kojima and Kangawa, 2005). The GHSR gene consists of 2 exons and 1 intron (Niarami *et al.*, 2014). In another study, the GHSR gene was shown to affect the fatty properties of meat in Xinghua chickens. GHSR is also related to the level of obesity in humans and mice. GHSR interacts with GHRL in the process of bioactivation through the neuroendocrine pathway (Lei *et al.*, 2007). According to Nie *et al.* (2009), the GHSR gene is also involved in GH secretion, food intake, and energy homeostasis. Fang *et al.* (2010) found a diversity of GHSR gene loci in intron 1 in White Recessive Rock and Xinghua chickens, which was associated with weights at 28 and 90 days of age, chest and thigh muscle weights. This diversity is caused by a mutation of the base T to C.

This study was conducted to investigate the diversity of GHSR gene genotypes in Bangkok chickens and the association of these genes with body weight, carcass weight, and commercial weight in Bangkok chickens.

## MATERIALS AND METHODS

### Ethical Approval

This study was approved by the Research Ethics Commission of Experimental Animals and

Health, Politeknik Negeri Jember, Number: 298/PL17.4/PG/2024.

### Study Period and Location

This study was conducted in July–December 2022. The locations of this study were the Teaching Laboratory, Department of Animal Science, Politeknik Negeri Jember, as a place of object cultivation, and the Genetic Molecular Laboratory, Faculty of Animal Science, IPB University, for the genetic observations.

### Livestock Samples

The sample used in this study was DOC Bangkok chickens, totaling 125 roosters, kept in colony cages. From the chicken population, 125 blood samples were collected from Bangkok chickens raised from DOC until 12 weeks. Food and drink were given *adlibitum*.

### Genotype Analysis

The total genome was obtained by the phenol-chloroform method (Sambrook and Russell, 2001). The use of primers to amplify the GHSR gene is referred to (Fang *et al.*, 2010) as modified. Forward primer (5'-CCC ACA AAG TTA GCT GCA GAC-3') and reverse primer (5'-CAC CTC TCC ATC TGG CTC ATT-3'). The primer was able to amplify the 470 bp GHSR gene in intron 1 (Figure 1).




Figure 1. Genomic organization and polymorphism sites of the chicken GHSR gene.

DNA amplification was carried out in a total volume of 15  $\mu$ L consisting of 1  $\mu$ L DNA, 0.3  $\mu$ L primer, 7.5  $\mu$ L master mix, and 6.2  $\mu$ L distilled water. PCR was carried out at the initial denaturation at 94°C for 4 minutes, which was carried out for 1 cycle. denaturation at 94°C for 10 seconds, annealing at 60°C for 20 seconds, extension at 7°C for 30 seconds (35 cycles), and final extension at 72°C for 7 minutes.

Determination of the genotype on the GHSR gene using the RFLP method. 5  $\mu$ L of the product from the GHSR gene PCR process was cut using a 2  $\mu$ L restriction endonuclease mix consisting of 1  $\mu$ L dH<sub>2</sub>O, 0.7  $\mu$ L buffer, and 0.3  $\mu$ L cutting enzyme, then incubated for 16 hours at 37°C. The cutting enzyme or restriction enzyme used for the GHSR gene is Hin6I, which can recognize the G|CGC cut site. Electrophoresis was carried out on a 2% agarose gel with 0.5 TBE buffer (Tris Borat EDTA), which was supplied with an electric current of 100 volts for 40 minutes. The visualization of electrophoresis results was carried out using a UV transilluminator.

### Data Analysis

The data analyzed was in the form of body proportion indicators, which included data on body weight, carcass weight, and commercial cuts (breast, thigh, drumstick, wings, and lumbar). Sequencing data from the GHSR gene were analyzed using the Bioedit program (Hall, 1999). Single Nucleotide Polymorphism (SNP) identification was carried out using the Molecular Evolutionary Genetics Analysis 7 (MEGA 7) program with reference number AB095994.1 (Tamura *et al.*, 2011). Allele frequency values, genotype frequencies, Hardy-Weinberg equilibrium values, observational heterozygosity, and expected heterozygosity were calculated based on Allendorf *et al.* (2012) method. The association between the GHSR gene and the observed traits in Bangkok chickens was analyzed using the T-test.

## RESULTS AND DISCUSSION

### GHSR Gene Amplification

The amplification of the GHSR gene in Bangkok chicken was successfully carried out with an initial denaturation process at 94°C for 4 minutes for 1 cycle. denaturation at 94°C for 10 seconds, annealing at 60°C for 20 seconds, extension at 72°C for 30 seconds (35 cycles), and final extension at 72°C for 7 minutes. The GHSR gene amplification process produces a product of 470 bp (Figure 2). The results of this amplification were similar to the amplification

process of the GHSR gene in Kampung chickens in the study of Syaikhullah *et al.* (2017). The results of the amplification were then cut using the restriction enzyme Hin6I and produced 2 types of genotypes, namely CT and TT. RFLP results for the GHSR gene showed that the CT genotype had 3 bands of 470 bp, 354 bp, and 116 bp, while the TT genotype had 1 band of 470 bp (Figure 3). These results are quite different when compared to the results of Fang *et al.* (2010) where the GHSR gene was found in Chinese local chickens (Xinghua) with as many as 3 genotype variants, namely CT, TT, and CC.

The sequencing results of the GHSR gene in Bangkok chickens found a deletion at the primary position of 134 bp, which caused the loss of base C (Figure 4). The alignment of the GHSR gene in Bangkok chickens indicates a mutation, where the T base changes to C. This can be referred to as a transversion mutation, specifically, a change from a purine base to a pyrimidine. SNP transversion has a high chance of causing amino acid changes resulting in non-synonymous mutations (Allendorf *et al.*, 2012).

According to Fang *et al.* (2010), mutations in the GHSR gene were observed to occur in intron 1. The intron also contains DNA sequences that bind enhancers that might affect the transcription process. Introns also contain regulatory RNA sequences that can affect the translation process and mRNA stability (Syaikhullah *et al.*, 2017). When a mutation occurs, RNA can change the product of a gene. Mutations in this area can affect the function of gene activity. In the process of translation, introns are also involved in the process of regulating protein production activity. Mutations in this region have the potential to affect the amount of protein production, function, or gene expression (Perdew *et al.*, 2006).

### Diversity, Heterozygosity, and HW Equilibrium of the GHSR Gene


Genotype frequency indicates the ratio of the number of a genotype to a population by calculating the ratio between the number of certain genotypes in each population, while allele frequency is the ratio of an allele to the total




number of alleles in an SNP in a population (Noor, 2010).

Table 1 showed that the frequency of the TT genotype in the GHSR gene is greater than that of the CT genotype, which was 0.776, while the frequency of the CT genotype was 0.224. In the GHSR gene, the frequency of the T allele was

0.888, and the frequency of the C allele was 0.112. The single nucleotide polymorphism (SNP) of the GHSR gene was indicated to be polymorphic. If an SNP has an allele frequency value of  $\leq 0.99$  in a large population and  $\leq 0.95$  in a smaller population, it can be stated that the SNP is polymorphic (Allendorf *et al.*, 2012).



**Figure 2.** Results of GHSR gene amplification in Intron 1 (S1–7 = Bangkok chicken blood sample).



**Figure 3.** RFLP results of the GHSR gene restricted by *Hin*6I enzyme; M: 100 bp; CT: Genotype CT (470 bp, 354 bp, 116 bp); TT: Genotype TT (470 bp).

DNA Sequences	Translated Protein Sequences
Species/Abbrv	G: *****
1. GENBANK	TTTAAGATCATGGAAGTGATCTTATTCTCTGCATTAGCAAAG
2. Genotipe_CT	TTTAAGATCATGGAAGCGATCTTATTCT-TGCATTAGCAAAG
3. Genotipe_TT	TTTAAGATCATGGAAGCGATCTTATTCT-TGCATTAGGAAAAG
4. 1811706_25X_GHSR_GHSR_R	TTTAAGATCATGGAAGCGATCTTATTCT-TGCATTAGGAAAAG

**Figure 4.** Spot C base deletion at position 134 in Bangkok chicken based on reference with GenBank access number AB095994.1.

**Table 1.** Genotype frequency, allele frequency, heterozygosity, and Hardy-Weinberg balance at the GHSR locus

Gen	Genotype Frequency		Allele Frequency		Ho	He	$\chi^2$
	TT (n = 97)	CT (n = 28)	T	C			
GHSR	0.776	0.224	0.888	0.112	0.224	0.198	1.99*

\* not significant ( $p > 0.05$ ,  $\chi^2 (0.05; 1) < 3.84$ ).

**Table 2.** Association of GHSR Intron 1 gene with body weight and carcass weight

Parameters	Genotype	
	TT	CT
Body weight (g)	774.61 ± 86.15 <sup>a</sup>	613.36 ± 32.39 <sup>b</sup>
Carcass weight (g)	463.25 ± 78.79 <sup>a</sup>	353.94 ± 31.09 <sup>b</sup>
Carcass percentage (%)	59.81 ± 2.8	57.71 ± 3.2

<sup>a,b</sup> different superscripts in the same row indicate significant difference ( $p < 0.05$ ).

**Table 3.** Association of GHSR Intron 1 gene with commercial weight

Parameters commercial weight	Genotype	
	TT	CT
breast (g)	117.92 ± 21.81 <sup>a</sup>	84.53 ± 7.73 <sup>b</sup>
thigh (g)	84.47 ± 13.52 <sup>a</sup>	67.15 ± 4.94 <sup>b</sup>
drumstick (g)	82.17 ± 12.32 <sup>a</sup>	55.73 ± 4.51 <sup>b</sup>
wing (g)	77.15 ± 11.14 <sup>a</sup>	55.64 ± 5.96 <sup>b</sup>
lumbar (g)	119.46 ± 23.93 <sup>a</sup>	86.15 ± 16.84 <sup>b</sup>

<sup>a,b</sup> different superscripts in the same row indicate significant difference ( $p < 0.05$ ).

Allendorf *et al.* (2012) explained that the diversity of a gene can be used as an indicator in determining a breeding program. The gene diversity number can be used as a tool to determine selection measures if the population is heterogeneous and attempts at crossing if the population is homogeneous (Fikri and Purnama, 2020). The value of heterozygosity is an indicator of the genetic diversity of a population that can be used for selection programs. An SNP is indicated to have high diversity if the heterozygosity value is  $> 0.50$  (Allendorf *et al.*, 2012; Wardhana *et al.*, 2019). The Ho value from Table 1 showed less than 0.50, which indicates that the genotypic diversity of the Bangkok chicken population in this study was considered low. Based on Table 1, the observed heterozygosity value (Ho) was 0.224, while the expected heterozygosity value (He) was 0.198. This study indicates that in the Bangkok chicken population under examination, there were no inbreeding cases. A lower Ho value than He indicates inbreeding (Nassiry *et al.*, 2009).

### Association of the GHSR Gene with Body Proportion in Bangkok Chickens

The association of the GHSR gene with body weight and carcass weight was presented in Table 2. Bangkok chickens showed that the TT genotype had higher body weights and carcass weights compared to the CT genotype Bangkok chickens. However, the carcass percentage of the two genotypes did not show a significant difference at the  $p < 0.05$  level. This shows that the GHSR gene is associated with body weight and carcass weight in Bangkok chickens.

These results follow the study of Syaikhullah *et al.* (2017), who found that the TT genotype was more dominant than the CT genotype in the body weight and carcass weight parameters of Kampung chickens. The GHSR gene at location rs16675844 had a significant effect on the characteristics of feed consumption and feed conversion, while the GHSR gene at location rs14678932 showed a significant association with the characteristics of breast weight and feed consumption (Jin *et al.*, 2014; Kartikasari *et al.*, 2019). The GHSR gene in chickens is the ghrelin

receptor (GHRL), while the GHSR gene is a peptide hormone produced by the chicken proventriculus, which stimulates the release of growth hormone (GH) and food intake (El-Magd *et al.*, 2016; Wardhana *et al.*, 2021). The GHSR gene can affect the growth and weight of chicken carcasses because these genes have an important role in biological response processes such as cell development and proliferation (Tschöp *et al.*, 2000).

Commercial weight is also one of the indicators used to observe the body proportions of chickens. The association of the GHSR gene with commercial cut weight can be seen in Table 3.

The results of statistical analysis showed that there were significant differences between the two genotypes ( $p < 0.05$ ) in breast weight, upper thigh weight, lower thigh weight, wing weight, and lumbar weight in Bangkok chickens. The TT genotype has a greater weight than the CT genotype in all commercial weight parameters. This shows that the GHSR gene is associated with commercial weight in Bangkok chickens. GHSR has also been extensively studied in several domestic animals, and SNPs are related to productivity performance. GHSR gene mutations were significantly ( $p < 0.05$ ) associated with several fatty properties such as crude protein content, thigh muscle weight, and thickness of subcutaneous fat in chickens and ducks (Nie *et al.*, 2009) Diversity in the GHSR gene was also found in a study conducted by Kaczor *et al.* (2016). The GHSR gene affects weight and daily weight gain in broiler chickens. A study conducted by El-Magd *et al.* (2016) found another mutation in the GHSR gene in exon 2 (G244A), which was a non-synonymous mutation, where there was a change in the amino acid product from lysine to arginine.

## CONCLUSION

The results of this study suggest that the Growth Hormone Secretagogue Receptor (GHSR) gene restricted by the Hin6I enzyme in Bangkok chickens has 2 genotypes, namely TT and CT. The two genotypes were associated with body weight, carcass weight, and commercial

weight ( $p < 0.05$ ). The TT genotype had a higher weight value than the CT genotype for all observed parameters. The GHSR gene has the potential to be used as a genetic marker for the selection process based on body weight, carcass weight, and commercial weight traits.

## ACKNOWLEDGEMENTS

The authors express their gratitude to the Department of Animal Science, Politeknik Negeri Jember, and the Faculty of Animal Science, IPB University, for supporting this study.

## AUTHORS' CONTRIBUTIONS

GS: Conceptualization and drafted the manuscript. GS, RTH, and MA: Performed sample evaluation. GS and MA: Performed the statistical analysis and the preparation of tables and figures. All authors have read, reviewed, and approved the final manuscript.

## COMPETING INTERESTS

The authors declare that they have no competing interests.

## REFERENCES

- Allendorf, F. W., Luikart, G. H., & Aitken, S. N. (2012). Conservation and the genetics of populations. John Wiley & Sons.
- Badaruddin, R., Astuty, F., & Pagala, M. (2017). Performa Penetasan Telur Ayam Hasil Persilangan Ayam Bangkok Dengan Ayam Ras Petelur. *JITRO*, 4(2).
- El-Magd, M. A., Saleh, A. A., Abdel-Hamid, T. M., Saleh, R. M., & Afifi, M. A. (2016). Is really endogenous ghrelin a hunger signal in chickens? Association of GHSR SNPs with increase appetite, growth traits, expression and serum level of GHRL, and GH. *General and Comparative Endocrinology*, 237, 131–139.

- Fang, M., Nie, Q., Luo, C., Zhang, D., & Zhang, X. (2010). Associations of GHSR gene polymorphisms with chicken growth and carcass traits. *Molecular Biology Reports*, 37, 423–428.
- Fikri, F., & Purnama, M. T. E. (2020). Biosecurity Application of Small Scale Chicken Abattoir in Sidoarjo, East Java, Indonesia. *Systematic Reviews in Pharmacy*, 11(6), 226–229.
- Hall, T. A. (1999). BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium Series*, 41(41), 95–98.
- Hijriyanto, M. (2017). Pengaruh frekuensi penampungan semen terhadap kualitas spermatozoa pada Ayam Bangkok. *Jurnal Ilmiah Mahasiswa Veteriner*, 1(1).
- Jin, S., Chen, S., Li, H., Lu, Y., Xu, G., & Yang, N. (2014). Associations of polymorphisms in GHRL, GHSR, and IGF1R genes with feed efficiency in chickens. *Molecular Biology Reports*, 41, 3973–3979.
- Kaczor, U., Poltowicz, K., Kucharski, M., Sitarz, A. M., Nowak, J., Wojtysiak, D., & Zieba, D. A. (2016). Effect of ghrelin and leptin receptors genes polymorphisms on production results and physicochemical characteristics of *M. pectoralis superficialis* in broiler chickens. *Animal Production Science*, 57(1), 42–50.
- Kartikasari, A. M., Hamid, I. S., Purnama, M. T. E., Damayanti, R., Fikri, F., & Praja, R. N. (2019). Isolation and Identification of *Escherichia coli* as Bacterial Contamination in Broiler Chicken Meat in Poultry Slaughterhouse Lamongan District. *Jurnal Medik Veteriner*, 2(1), 66–71.
- Kojima, M., & Kangawa, K. (2005). Ghrelin: structure and function. *Physiological Reviews*, 85(2), 495–522.
- Lei, M., Luo, C., Peng, X., Fang, M., Nie, Q., Zhang, D., Yang, G., & Zhang, X. (2007). Polymorphism of growth-correlated genes associated with fatness and muscle fiber traits in chickens. *Poultry Science*, 86(5), 835–842.
- Muryanto, H. P., Herman, R., & Setijanto, H. (2002). Evaluasi Karkas Hasil Persilangan antara Ayam Kampung Jantan dengan Ayam Ras Petelur Betina. *Journal Animal Production*, 4(2), 71–76.
- Nassiry, M. R., Javanmard, A., & Reza, T. (2009). Application of statistical procedures for analysis of genetic diversity in domestic animal populations. *American Journal of Animal and Veterinary Sciences*, 4(4), 136–141.
- Nataamijaya, A. G. (2010). Pengembangan potensi ayam lokal untuk menunjang peningkatan kesejahteraan petani. *Jurnal Litbang Pertanian*, 29(4), 131–138.
- Niarami, D. M., Masoudi, A. A., & Vaez Torshizi, R. (2014). Association of single nucleotide polymorphism of GHSR and TGFB2 genes with growth and body composition traits in sire and dam lines of a broiler chicken. *Animal Biotechnology*, 25(1), 13–22.
- Nie, Q., Fang, M., Xie, L., Peng, X., Xu, H., Luo, C., Zhang, D., & Zhang, X. (2009). Molecular characterization of the ghrelin and ghrelin receptor genes and effects on fat deposition in chicken and duck. *BioMed Research International*, 2009.
- Ningsih, N., Syaikhullah, G., Adhyatma, M., Rofiqi, Z., & Al Huzaini, M. F. (2022). The effect of phytobiotics ginger (*Zingiber officinale* Rosc.) extract in drinking water on hematological characters of KUB chickens. *IOP Conference Series: Earth and Environmental Science*, 980(1).

- Perdew, G. H., Vanden Heuvel, J. P., & Peters, J. M. (2006). Regulation of gene expression: molecular mechanisms. Springer.
- Ronny Rachman Noor. (2010). Genetika Ternak. Penebar Swadaya.
- Sambrook, J., & Russell, D. W. (2001). Molecular Cloning: Ch. 8. In Vitro amplification of DNA by the polymerase chain reaction (Vol. 2). Cold Spring Harbor Laboratory Press.
- Syaikhullah, G., Darwati, S., & Sumantri, C. (2017). Asosiasi Keragaman Growth Hormone Secretagogue Receptor (GHSR| Hin6i) terhadap Bobot Karkas Ayam Kampung pada Umur 12 Minggu. *Jurnal Ilmu Produksi Dan Teknologi Hasil Peternakan*, 5(3), 117–121.
- Tschöp, M., Smiley, D. L., & Heiman, M. L. (2000). Ghrelin induces adiposity in rodents. *Nature*, 407(6806), 908–913.
- Wardhana, D. K., Haskito, A. E. P., Purnama, M. T. E., Safitri, D. A., & Annisa, S. (2021). Detection of microbial contamination in chicken meat from local markets in Surabaya, East Java, Indonesia. *Veterinary World*, 14(12), 3138–3143.
- Wardhana, D. K., Purnama, M. T. E., Kean, O. H., & Tyasningsih, W. (2019). Detection of salmonella on chicken meat using immunomagnetic separation and conventional methods from traditional market in Surabaya, East Java, Indonesia. *Indian Veterinary Journal*, 96(11), 31–33.

\*\*\*