

DAFTAR PUSTAKA

- Acar, B., Sancaktutar, A. A., & Bozkurt, H. (2021). "Effect of obesity on percutaneous nephrolithotomy outcomes in Staghorn stones." *New Journal of Urology*, 14(1): 16-21.
- Chen, B., Wang, M., Chen, Z., Pan, N., He, X., Li, B., Chen, C., Zhou, Z., Cui, W., Zhang, P., Ren, Q., & Zhong, Y. (2024). Clinical comparison of lateral-supine mini-percutaneous nephrolithotomy and anatrophic nephrolithotomy in the treatment of complete staghorn calculi. *BMC Urology*, 24, 167.
- Keheila, M., et al. (2023). "Efficacy and safety of percutaneous nephrolithotomy in patients with body mass index over 30 kg/m²." *Journal of Endoscopic and Minimally Invasive Surgery*, 3(2): 113-118.
- Kementerian Kesehatan RI. (2014). *Buku Pedoman Proses Asuhan Gizi Terstandar*. Jakarta: Kementerian Kesehatan RI.
- Martínez-Corral, M. E., Vázquez-Cancela, O., Fernández-Pérez, C., & Pérez-Fentes, D. A. (2025). Impact of Obesity on the Outcomes of Supine Percutaneous Nephrolithotomy: A Propensity Score Analysis. *Journal of Endourology*, 39(3), 237-244.
- Xu, Y., & Huang, X. (2022). Effect of body mass index on outcomes of percutaneous nephrolithotomy: a systematic review and meta-analysis. *Frontiers in Surgery*, 9, 922451.