

## DAFTAR PUSTAKA

- Aaqil, M., Peng, C., Kamal, A., Nawaz, T., Zhang, F., & Gong, J. (2023). Tea Harvesting and Processing Techniques and Its Effect on Phytochemical Profile and Final Quality of Black Tea: A Review. *MDPI*, *12*, 1–28. <https://doi.org/https://doi.org/10.3390/foods12244467>
- Ashihara, H. (2015). Occurrence, biosynthesis and metabolism of theanine ( $\gamma$ -glutamyl-L-ethylamide) in plants: a comprehensive review. *Natural Product Communications*, *10*(5), 803–810.
- Badan Pusat Statistik, B.-S. I. (2024). *Statistik Teh Indonesia* (Vol. 17).
- Bilardi, P. A. (2023). Implementasi HACCP (Hazard Analysis and Critical Control Point) pada Proses Produksi Teh Sencha di PT. Bali Cahaya Amerta Denpasar. *FTP-UB e-Library*. [https://ftp-e-library.ub.ac.id/koleksi\\_detail.php?backurl=koleksi\\_kategori.php@kodekat%3D002.01&kode\\_koleksi=310012300014](https://ftp-e-library.ub.ac.id/koleksi_detail.php?backurl=koleksi_kategori.php@kodekat%3D002.01&kode_koleksi=310012300014)
- Collings, E. R., Alamar, M. C., Redfern, S., Cools, K., & Terry, L. A. (2019). *Spatial changes in leaf biochemical pro fi le of two tea cultivars following cold storage under two di ff erent vapour pressure de fi cit ( VPD ) conditions*. *277*(July 2018), 179–185.
- Cronquist, A. (1981). *An Integrated System of Classification of Flowering Plants*. Columbia University Press.
- Effendi, D. S., Syakir, M., Yusron, M., & Hartati, R. S. (2017). *Budidaya dan Pascapanen Teh*. IAARD Press.
- Fitzpatrick, J. (2013). Powder properties in food production systems. *Woodhead Publishing*, 285–308. <https://doi.org/https://doi.org/10.1533/9780857098672.2.285>.
- Ishiguro, Y., Yamashita, H., & Ikka, T. (2026). Advances in tea breeding in Japan : from traditional methods to genomic approaches. *Japan Agricultural Research Quarterly (JARQ)*, *53*(1), 1–11. <https://doi.org/10.1270/jsbbs.25050>
- Kohata, K., Yamashita, Y., Yamauchi, Y., & Horie, H. (1999). Investigations of Qualities and Properties of Commercially Available Green Teas on the Basis of Chlorophyll and its Derivatives Contents and Hideki HORIE National of

Vegetables , Ornamental Plants and Tea. *J-Stage*, 1999(87), 13–19.  
<https://doi.org/https://doi.org/10.5979/cha.1999.13>

Liu, Z., Ran, Q., Li, Q., Yang, T., Dai, Y., Zhang, T., Fang, S., Pan, K., & Long, L. (2023). Interaction between major catechins and umami amino acids in green tea based on electronic tongue technology. *Food Science*, 88(6), 2339–2352.  
<https://doi.org/10.1111/1750-3841.16543>

Matsuda, T., Matsugatani, Y., & Hara, M. (2016). *A New Shading Method for Improving Free Amino Acid Content and Yield of ‘ Yabukita ’ Kabuse-cha*. 9–15.

Ni, R., Yan, H., Tian, H., Zhan, P., & Zhang, Y. (2022). Characterization of key odorants in fried red and green huajiao (*Zanthoxylum bungeanum* maxim. and *Zanthoxylum schinifolium* sieb. et Zucc.) oils. *Food Chemistry*, 377.  
<https://doi.org/https://doi.org/10.1016/j.foodchem.2021.131984>.

Qin, W., Yamada, R., Takuya, A., & Ogawa, Y. (2022). Changes in Morphological and Functional Characteristics of Tea Leaves During Japanese Green Tea (Sencha) Manufacturing Process. *Food and Bioprocess Technology*, 15, 82–91. <https://doi.org/10.1007/s11947-021-02735-7>

Rodiana, D. (2015). *Teh: Proses, Karakteristik & Komponen Fungsionalnya* (Vol. 10, Issue 8).